



Psychometric properties of histrionic personality symptoms scale among Iranian students

Sohrab Amiri¹, Yusef Jamali²

Journal of Research & Health
Social Development & Health Promotion
Research Center
Early View 26 Jun 2019
Original Article

1. **Correspondence to:** Behavioral Sciences Research Center, Lifestyle Institute, Baqiyatallah University of Medical Sciences, Tehran, Iran

Email: Amirysohrab@yahoo.com

2. Department of Counseling, Faculty of Psychology and Educational Sciences, Isfahan University, Isfahan, Iran

Received: 30 Jan 2017

Accepted: 15 Apr 2018

How to cite this article: Amiri S, Jamali Y. Psychometric properties of histrionic personality symptoms scale among Iranian students. *J Research & Health* 26 Jun 2019; Early View.

Abstract

Hysterical personality is a personality disorder in the international classification of disease and diagnostic and statistical manual of mental disorders. This study was aimed to evaluate the psychometric properties of histrionic personality symptoms scale. The study was methodology. For this purpose, 322 students were selected by using multi-stage cluster sampling then histrionic personality scale, histrionic personality subscale from millon clinical multiaxial inventory and the big five-factor personality inventory were distributed among them. After collecting data, initially the questionnaire was translated into Persian with back translation techniques to determine the psychometric properties of histrionic personality symptoms scale and validity of the scale was assessed by Cronbach's alpha, retest and splitting coefficient, which were 77.0, 89.0, 56.0, 49.0, 67.0, 71.0 that the criterion validity with histrionic personality subscale from millon clinical multiaxial inventory and five-factor personality inventory. As well as its factor structure was assessed by using exploratory and confirmatory factor analysis, fit indexes of X^2/df equal to 3.2, SRMR=0.07, CFI=0.90 and RMSEA=0.10. The results of the factor analysis indicated that the histrionic personality scale has 2 factors, test the reliability of the scale by using Cronbach's alpha, retest and splitting coefficient reflects the reliability of the scale, the criterion validity of the questionnaire with other questionnaires showed desirable discriminant and convergence validity. In Overall, these findings indicated histrionic personality scale has good psychometric properties among Iranian populatoin.

Keywords: Histrionic, Personality, Reliability, Scale, Validity

Introduction

The hysterical personality has been showed in both the international classification of diseases and health problems [1] and the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) [2]. Hysterical Personality Disorder (HPD) is a clinical syndrome in which people have an interactive style interpersonal style which is characterized by seduction, emotional superficiality, and displaying by seductiveness,

emotional superficiality, and dramatics [2]. According description in the DSM-5; [2], people with HPD have being motivated by a search for being the center of attention; these are seductive, flirting and excite other people's sex, high self-expression in expressing emotions also has an emotional superficiality and ignores the others feelings that these people often describe themselves as attractive

people and may perform sexually suggestive behaviors. People with hysterical personalities often have high social and occupational functioning, despite being easily influenced by others and tend to use their skills to control and manage others but showing good social skills [3-5]. Because of the nature of being dramatic and extraversion, people with HPD may have some fans, although they often have problems in maintaining a deep and mutually satisfactory relationship. The prevalence of this disorder in the general population is estimated at 2-3% [6]. This disorder is often diagnosed in women [4], although some studies have reported the equal prevalence in both genders [7].

Kraus and Reynolds [6] stated that hysterical personality disorder was associated with an increased risk for a variety of disorders, including depression and anxiety and argue that HPD can be an important clinical syndrome for study. It has been shown that hysterical personality is associated with a number of negative consequences including reduced marital satisfaction and success [8] and higher rates of depression disorder [9]. Compared to other Class B personality disorders, few researches have been focused on HPD. Although there may be several reasons that have been less addressed to HPD that one reason may be due to the lack of a HPD symptom scale is available and at the same time concise, short, and valid. Clinical Multidimensional Millon Clinical Questionnaire (MCMI) [10] is the only clinical test involving a hysterical scale. On the contrary, the Minnesota Personality Inventory [11] does not include any scale from the HPD, although hysterical personality traits may be inferred from scales. Colligan, Morey and Offord [12] developed a prestigious hysterical scale for MMPI. However, due to the lack of scoring pattern this scale is inappropriate for clinicians. Both the MCMI-III and the MMPI-2 are long because of their large personality scales and its scoring on the hysterical scale of MMPI-2 is inappropriate. In this way, it seems that a clinical measure scale for HPD traits that is concise, valid, accessible, and easy to scoring for clinical and research populations is

necessary. Accordingly, Ferguson and Noguee [13] formulated a screening scale to measure hysterical personality symptoms. This scale consists of 11 questions and two subscales of seductiveness and attention seeking which is standardized on two samples of 661 and 340 adults that results have shown this scale has favorable psychometric properties and hence it can be used as a valid tool to measure hysterical personality traits. As above, despite the fact that hysterical personality traits have clinical significance, however the study of these traits due to the lack of appropriate tools doesn't have enough research and clinical attention so the purpose of this study was to evaluate the psychometric properties Hysterical Personality Symptom Scale (BHPS) among Iranian university students.

Method

The population of the study was all students of Bu-Ali Sina university in 2016-2017 academic years. The participants of the study were 322 students who were selected by multistage cluster sampling among faculties (full-time, tuition-based program and part-time). Data were gathered using survey method (questionnaire).

Hysterical Personality Symptom Scale (BHPS): Initially, in order to shape the hysterical personality scale, a set of 36 items was selected that had DSM-5 criteria for the identification of hysterical personality disorder. Questions should be responded based on the 4-level Likert spectrums are never to ever. After exploratory factor analysis using maximum likelihood method with promax rotation 11 times with higher 0.60 factor load was extracted. Based on this, factor analysis yielded two-factor solution. 11 items had alpha coefficients of 0.76. Seductiveness subscale has been shown alpha coefficient of 0.67 and subscale of attention seeking 0.74 [13]. In order to prepare the Persian version of the BHPS [13], the researcher and two English language experts translated the questionnaire to Persian then the back translated text was revised. In the next step, the translated text, translated into

English by the researcher and English language experts and the two versions were matched to each other. Then, the questionnaire was given two psychology experts to test its Facial Validity. After the match between the translated version and the original and solving the defects, questionnaire was implemented on 40 students similar to the final research community and then getting the feedback about the content of the items and fixing the weaknesses the final version of the questionnaire was prepared for use.

Multidimensional Millon Clinical Questionnaire-3 (MCMI-III): A self-assessment scale with 175 items which evaluates 14 clinical patterns of personality and 10 clinical syndromes and is used for adults aged 18 or over for therapeutic or psychological evaluation. Its validity has been confirmed through factor analysis and its reliability has been confirmed through internal consistency and proper test-retest. The retest reliability coefficient was reported at a median of 0.91. The range of retest coefficients was between 5 and 14 days [14]. Khajeh Mughni [15] conducted a study in order to standardize this test in Tehran. In this study, the reliability coefficient of MCMI scales was obtained from a range of 7 to 10 days from a range of 0.78 (dramatic scale) to 0.87 (delusional disorder scale).

Neo Personality Questionnaire (NEO): This questionnaire is a sixty-item that can measure five main factors and six attributes of each factor. In this form, participants receive a score of 0-4 in each sentence and every twelve questions are one of the factors of scale [16]. McCrae and Costa in 1992 reported Cronbach's

Alpha coefficients of 0.66 for agreeableness to 0.86 for Neuroticism [16]. This test has been implemented and standardized in Iran by Hagnosh, whose desirable Cronbach's alpha coefficients have been reported in 512 participants [17].

In order to collect data, the researcher first described the response process and then distributed tools to answer among the participants. During the research process, the researcher was present among the participants in order to ensure the accuracy of the response, as well as to resolve the ambiguity. Data were analyzed by SPSS-22 and LISREL 8.8 [18]. Psychometric indicators of the BHPS were calculated through descriptive indexes, reliability and validity coefficients, confirmatory factor analysis.

Results

Of the participants, the number of undergraduate students was 194 (60.3%), the master degree was 106 (32.9%) and 22 (6.8%) were PhD. The descriptive and demographic characteristics of the participants showed that 65 boys (20.2%), 257 girls (79.8%), as well as means and standard deviations was obtained for boys age 72.21 (30.2) and for girls 29.21 (18.2) respectively.

Validity of Persian version of the BHPS [13] was investigated with two methods of critical validity Simultaneous implementation with hysterical personality subscale from Millon clinical multiaxial inventory [14] and The big five-factor personality inventory [16], and the correlation between subscales (Table 1).

Table 1 Correlation coefficients between subscales of hysterical personality signs with other scales

Subscales	1	2	Hysteric (million)	Extroversion	Conscientiousness	Agreeableness	Neuroticism	Openness
Seductiveness	1	-	0.38**	0.29**	-0.08	0.07	0.11*	0.13**
Attention seeking	0.54**	1	0.36**	0.42**	-0.10	0.09	0.06	0.27**

**p<0.01 *p<0.05

The pattern of correlation coefficients between subscales in Table 1 shows that there is a proper internal relation between subscales. Also, the correlation coefficient model of the subscales with the hysterical personality subscale from

Millon vlinical multiaxial inventory [14] and The big five-factor personality inventory [16] indicates the simultaneous validity of the BHPS scale [13].

In order to investigate the fitting of the two-

factor structure of the Persian version of the BHPS [13], a confirmatory factor analysis using maximum likelihood method and the LISREL software was used [18].

The path diagram of the confirmatory factor analysis along with the path coefficients in Figure 1 and also the T index in Table 2 has been presented.

Table 2 T Index for evaluation of significant level

Subscales	ITEM	T index	Subscales	ITEM	T index
Seductiveness	1	11.75	Attention seeking	2	11.35
	4	2.61		3	6.44
	5	4.73		6	7.53
	9	6.28		7	11.55
	10	7.84		8	10.09
	11	9.73			




Figure 1 Confirmatory factor analysis graph and path coefficients of Hysterical personality symptom scale

Table 3 Fitness indices of Hysterical personality symptom scale

Index	X ²	df	Value	RMSEA	SRMR	NFI	NNFI	RFI	IFI	GFI	AGFI	CFI
Value	138.50	43	<0.05	0.10	0.07	0.80	0.80	0.76	0.84	0.90	0.84	0.84

The confirmatory factor analysis by using LISREL software offered 3 levels of fit indices [19,20]: 1) absolute fit indices such as Chi-square index and Standardized Root Mean Square Residual (SRMR), 2) parsimony goodness-of-fit index such as Root Mean Square Error of Approximation (RMSEA) and 3) Comparative Fit Index (CFI). There is controversy over the precise cutting scores of fit indices [19-23]. More the scores of Normed Fit Index (NFI), Relative Fit Index

(RFI), and Goodness of Fit Index (GFI) are closer to 1; they indicate the pattern's more desired fit. In the present study, the most valid fit scores have been used in order to assess the model's fit. SRMR<0.08 shows a desired fit and SRMR<0.10 indicates an acceptable fit and the model is failed when SRMR>.10 [22]. The index with value of RMSEA≤.8 shows the model is good, and when the RMSEA is between 0.10 and 0.08, the model is acceptable. Eventually, CFI≥.95

states the good fit of the model [22]. Since the X^2 statistic is sensitive to the sample size, to assess the overall fit of the model, the amount of X^2 is calculated along with the degree of freedom (X^2/df). $X^2/df < 2$ represent the model's good fit and when X^2/df is around 3, the model is acceptable. The amount of X^2/df was 3.2, which is around 3, so the model's fit is acceptable. In addition to this, SRMR=.07 showed the 6 factor model's fit is acceptable and the fit indices of CFI=.90 and RMSEA=.10 represent the model's acceptable fit.

To assess the reliability of the BHPS [13], the Cronbach's Alpha coefficient, Two-half coefficient and test-retest were calculated. The Cronbach's alpha coefficient of 11 questions was 0.75, indicating that the scale had good internal consistency. Also, two-half coefficients

of the total scale 0.77 and its subscales 0.67 and 0.71 showed good consistency of the scale and its sub-scales.

In order to calculate the retest coefficient, 47 participants responding to the questionnaire two weeks later and correlation coefficient between two times the implementation of the questionnaire was calculated. The results of the Cronbach's alpha coefficient, the two-half and test-retest coefficient would be presented in Table 4.

The results of Table 4 show that the Cronbach's alpha coefficients are satisfactory and also the data of all the coefficients of the retest and the two half-sets are significant. Therefore, it can be concluded that the Persian version of the BHPS [13] has a favorable internal consistency.

Table 4 Mean, standard deviation, alpha coefficients, and Hysterical personality symptoms subscales

Subscale	ITEM	M/SD	Alpha coefficient	Retest coefficients (N = 47)	Two-half coefficient
Seductiveness	1, 4, 5, 9, 10, 11	12.11 (3.03)	0.77	0.56**	0.67
Attention seeking	2, 3, 6, 7, 8	12.48 (2.98)	0.89	0.49**	0.71

** $p < 0.01$

Discussion

The present study evaluates the structure and validity of the BHPS by examining the reliability, validity and factor structure. The present findings provide clear support for the reliability and validity of the BHPS as new questionnaire to eliminate the limitations of existing questionnaires and valid measurement of hysterical personality traits. As expected from previous studies [13], factor analysis of BHPS questionnaire showed that the two-factor solution has a desirable fit. This finding is consistent with studies which examined the factor structure of the BHPS [13]. In addition, the results showed that two sub-scales include: 1) Seductiveness 2) Attention seeking has a favorable internal reliability. Investigating the factor structure and factor load patterns of the BHPS scale, using the confirmatory factor analysis, obtained similar results to Ferguson and Negy [13] two-factor solution. Therefore, the findings of this study are consistent with the results of Ferguson and Negy's research [13] that shows the BHPS questionnaire has a

desirable structure.

The study of the reliability of the BHPS, by calculating the alpha coefficients, the coefficients of test-retest and Two-half coefficient indicated a desirable reliability of this scale. Alpha coefficients of subscales: 1) Seductiveness 2) Attention seeking were 0.77 and 0.89, respectively. And the re-test and the two-half coefficient indicate the desirability of the reliability of the BHPS; these findings are in line with the study of Ferguson and Negy [13] research which formed the original version of the BHPS.

The validity of the hysterical personality trait scale with hysterical personality subscale of Millon Clinical Multiaxial Inventory [14] shows positive correlation coefficients among all the similar subscales in the hysterical personality scale and the hysterical personality subscale ($p < 0.01$) this result demonstrates the convergence validity of the BHPS.

Psychometric properties of hysterical

personality symptoms scale in the present study are consistent with studies in the original version [13]. According above, the lack of a specific and useful instrument for measuring hysterical personality dimensions is one of the weaknesses in the field of personality research. Consequently, regardless of language and culture, and considering the pattern of factor loads consistent with previous studies [13], the hysterical personality scale seems to be a useful tool for better understanding the hysterical fundamental constituents of the personality. Generally, reliability, validity and confirmatory factor analysis revealed the desirable psychometric characteristics of hysterical personality symptoms scale and the findings of the present study are consistent with the results of the original version [13] and that the Persian version of the hysterical personality symptom showed that it was a reliable tool for assessing the dimensions of hysterical traits. Also, the calculated indices for the fitting of the hysterical personality marker scale model showed that SRMR (RMSEA), (X^2/df) as the most prestigious fitness indicators [19-23] support the fitting of the model. This finding was in line with the results of the original version of the BHPS [13]. So according to what was said, the present study was conducted among the student population. It is also recommended that in future research; the validity of the BHPS should be studied using other clinical approaches. Although the present study mainly focused on hysterical personality symptoms scale from the self-report perspective, however, adding information from clinical practices could be attractive and provide more information. On the other hand, information was gathered in a cross-sectional manner, thus limiting the conclusion about the observed effects. So future studies should have focus on longitudinal research. In total the result of this study showed that the validity and reliability of the hysterical personality trait scale in the sample of normal Iranian population. In addition, the factor structure of the questionnaire was consistent with the theoretical view of its makers and the two factors extracted based on the confirmatory

factor analysis were in line with the original version of the BHPS. Despite the limitations expressed, the present findings show that the BHPS can measure the components of hysterical personality and distinguish between their dimensions.

Conclusion

In total, hysterical personality symptoms scale has the good psychometric properties and applicability in the wide range and measurement of hysterical traits it also has the ability to be used at research, clinical, and personality levels.

Acknowledgments

The authors of the article need to acknowledge the cooperation and assistance of all the loved ones who helped us in this research.

Authors' contributions

Study design: AS

Data collection and analysis: AS, YJ

Manuscript preparation: AS, YJ

All authors have read and approved the final version

Conflict of Interest

"The authors declare that they have no competing interests."

Funding

The author (s) received no financial support for the research, authorship and/or publication of this article.

Availability of data and materials

The datasets used and/or analyzed during this study are available from the corresponding author on reasonable request.

References

- 1- World Health Organization. The ICD-10 classifications of mental and behavioral disorders: Clinical descriptions and diagnostic guidelines. Geneva: World Health Organization; 1992.
 - 2-American Psychiatric Association. Diagnostic and statistical manual of mental disorders—5. Washington, DC: American Psychiatric Association; 2013.
-

- 3- Weiner IB, Craighead WE. The corsini encyclopedia of psychology. Hoboken, New Jersey, US: John Wiley & Sons; 2010.
- 4- Millon T, Davis R. Disorders of personality: DSM-IV and beyond, 2nd ed. Oxford, England: John Wiley & Sons; 1996.
- 5- Shedler J, Westen D. Refining personality disorder diagnosis: integrating science and practice. *Am J Psychiatry* 2004; 161(8), 1350-65.
- 6- Kraus G, Reynolds D. The "A-B-C's" of the cluster B's: identifying, understanding, and treating cluster B personality disorders. *Clin Psychol Rev* 2001; 21(3): 345-73.
- 7- Nestadt G, Romanoski A, Chahal R, Merchant A, Folstein MF, Gruenberg EM, McHugh PR. An epidemiological study of histrionic personality disorder. *Psychol Med* 1990; 20(2): 413-22.
- 8- Disney KL, Weinstein Y, Oltmanns TF. Personality disorder symptoms are differentially related to divorce frequency. *J Fam Psychol* 2012; 26(6), 959-65.
- 9- Bockian NR. Depression in histrionic personality disorder. In *Personality guided therapy for depression* (pp. 169-186). Washington, DC, US: American Psychological Association; 2006.
- 10- Millon T, Millon C, Davis RD, Grossman S. Millon clinical multi-axial inventory-III (MCMI-III). London, UK: Pearson; 2009.
- 11- Hathaway S, McKinley J. MMPI-2: Manual for administration and scoring. Minneapolis: University of Minnesota Press; 1989.
- 12- Colligan R, Morey L, Offord K. The MMPI/MMPI-2 personality disorder scales: Contemporary norms for adults and adolescents. *J Clin Psychol* 1994; 50(2): 168-200.
- 13- Ferguson CJ, Negy C. Development of a brief screening questionnaire for histrionic personality symptoms. *Pers Individ Dif* 2014; 66, 124-7.
- 14- Marnat GG. Handbook of psychological assessment. Hoboken, New Jersey, US: John Wiley & Sons; 2003.
- 15- Khajeh Moogi N. Preliminary preparation of the persian form of Millon Clinical Questionnaire-2 in Tehran. [thesis]. Clinical psychology. Tehran: Psychiatric Institute, Iran University of Medical Sciences 1993; pp: 71-82.
- 16- Costa PT, McCrae RR. Revised NEO Personality (NEO-PI-R) and NEO Five-factor Inventory (NEO-FFI) Professional manual. Odessa, Florida, US: Psychological assessment resources; 1992.
- 17- Haq Shenaz H. Personality psychology. Shiraz, Iran: Shiraz university press; 2009. pp: 40-8.
- 18- Jöreskog KG, Sörbom D. LISREL 8.80 for windows (Computer Software). Lincolnwood, IL: Scientific Software International, Inc; 2006.
- 19- Brown TA. Confirmatory factor analysis for applied research. New York, US: Guilford publications; 2014
- 20- Schermelleh-Engel K, Moosbrugger H, Müller H. Evaluating the fit of structural equation models: Tests of significance and descriptive goodness of fit measures. *Psychol Res* 2003; 8(2): 23-74.
- 21- Hair JF, Black WC, Babin BJ, Anderson RE. Multivariate data analysis. Upper Saddle River, NJ: Pearson prentice hall; 2006.
- 22- Hu LT, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: conventional criteria versus new alternatives. *Structural Equation Modeling A: Multidisciplinary Journal* 1999; 6(1): 1-55.
- 23- Tabachnick BG, Fidell LS. Using multivariate statistics, 5th ed. Boston, MA: Pearson; 2007.